



## 第 3 次作业

危国锐 516021910080

(上海交通大学电子信息与电气工程学院, 上海 200240)

**摘要:** 第三章作业. 截止日期: 2022-03-25.

**关键词:** 关键词 1, 关键词 2

## Homework 3

Guorui Wei 516021910080

(School of Electronic Information and Electrical Engineering,  
Shanghai Jiao Tong University, Shanghai 200240, China)

**Abstract:** Selected exercises for chapter 3. Due date: Mar. 25<sup>th</sup>, 2022.

**Keywords:** keyword 1, keyword 2



# 目 录

<b>摘要 .....</b>	<b>i</b>
<b>Abstract.....</b>	<b>i</b>
<b>1 Question 2 .....</b>	<b>1</b>
1.1 Solution.....	1
1.1.1 Brief Review: <i>Lambert–Bouguer–Beer Law of Extinction</i> .....	1
1.1.2 A Special Case: Isothermal Plane-parallel Atmosphere in Hydrostatic Balance .....	1
<b>2 Question 1 .....</b>	<b>2</b>
2.1 Solution.....	2
<b>References .....</b>	<b>3</b>



# 1 Question 2

“Derive (3.23) using (3.22), (3.19) and the ideal gas law.”(Hartmann, 2016, p. 93)

## 1.1 Solution

除特别说明外，本节各符号的含义同 Hartmann (2016) 中的.

### 1.1.1 Brief Review: Lambert–Bouguer–Beer Law of Extinction

由 Lambert–Bouguer–Beer 消光定律 (*law of extinction*)

$$dF = -F \rho M_a k_{\text{abs}} ds, \quad ds = -\sec \theta dz \quad (1.1)$$

得

$$\frac{dF}{F} = \rho M_a k_{\text{abs}} \sec \theta dz. \quad (1.2)$$

对  $z$  从高度  $z = z$  到大气上界  $z = +\infty$  积分，得

$$F = F_\infty \exp(-\tau \sec \theta), \quad (1.3)$$

其中  $F_\infty, \theta$  是常数，

$$\tau := \int_z^{+\infty} \rho M_a k_{\text{abs}} dz \quad (1.4)$$

被称为沿传输路径的光学厚度 (*optical depth*). 由 (1.3) 得

$$dF = -F \sec \theta d\tau, \quad (1.5)$$

从而吸收率 (*Absorption rate*)

$$\frac{dF}{dz} = -F_\infty \exp(-\tau \sec \theta) \sec \theta \frac{d\tau}{dz}. \quad (1.6)$$

下面将本节的结果应用于静力平衡状态下的平面平行等温大气.

### 1.1.2 A Special Case: Isothermal Plane-parallel Atmosphere in Hydrostatic Balance

将理想气体状态方程

$$p = \rho RT \quad (1.7)$$

代入静力平衡方程

$$dp = -\rho g dz \quad (1.8)$$

得

$$\frac{dp}{p} = \frac{d\rho}{\rho} = -\frac{dz}{H}, \quad (1.9)$$

其中

$$H := RT/g \quad (1.10)$$

为大气标高 (scale height). 对 (1.10) 从下表面 ( $z = 0$ ) 到高度  $z = z$  积分，得

$$p = p_s \exp(-z/H), \quad (1.11)$$

$$\rho = \rho_s \exp(-z/H). \quad (1.12)$$

将 (1.12) 代入 (1.4)，假定  $M_a, k_{\text{abs}}$  为常数，得

$$\tau = k_{\text{abs}} M_a \rho_s H \exp(-z/H), \quad (1.13)$$

从而



$$d\tau = -\tau dz/H. \quad (1.14)$$

将 (1.14) 代入 (1.6), 得

$$\frac{dF}{dz} = \frac{F_\infty}{H} \tau \exp(-\tau \sec \theta) \sec \theta. \quad (1.15)$$

易知吸收率  $dF/dz$  (作为  $\tau$  的函数) 有最大值, 下求之.

令

$$0 = \frac{d}{d\tau} \ln \frac{dF}{dz} = \frac{1}{\tau} - \sec \theta, \quad (1.16)$$

得  $\ln(dF/dz)$  的唯一驻点

$$\tau = \tau_{\max \text{abs}} := \cos \theta, \quad (1.17)$$

故吸收率  $dF/dz$  在  $\tau = \cos \theta$  处取最大值. 将 (1.17) 代入 (1.13) 得该最大值点位于高度

$$z = z_{\max \text{abs}} := H \ln(k_{\text{abs}} M_a \rho_s H \sec \theta) \quad (1.18)$$

处, 再由 (1.11) 得该处的压强为

$$p = p_{\max \text{abs}} := p_s \frac{\cos \theta}{H k_{\text{abs}} \rho_{\text{as}}}, \quad (1.19)$$

其中

$$\rho_{\text{as}} = M_a \rho_s \quad (1.20)$$

“is the density of the absorber at the surface.” ([Hartmann, 2016, p. 64](#))

## 2 Question 1

“Suppose a gas that absorbs solar radiation has a uniform mixing ratio of  $1 \text{ g kg}^{-1}$  and an absorption cross-section of  $5 \text{ m}^2 \text{ kg}^{-1}$ . At what altitude will the maximum rate of energy absorption per unit volume occur? Assume an isothermal atmosphere with  $T = 260 \text{ K}$ , and a surface pressure of  $1.025 \times 10^5 \text{ Pa}$ , and that the Sun is directly overhead.” ([Hartmann, 2016, p. 93](#))

### 2.1 Solution

将 (1.7) (1.10) 代入 (1.18) 得

$$z = z_{\max \text{abs}} := \frac{RT}{g} \ln \left( k_{\text{abs}} M_a \frac{p_s}{g} \sec \theta \right), \quad (2.1)$$

代入  $\theta = 0$  和其他数据得

$$z_{\max \text{abs}} = 30091 \text{ (m)}. \quad (2.2)$$



## References

- Hartmann, D. L. (2016). Chapter 3 - Atmospheric Radiative Transfer and Climate. In D. L. Hartmann (Ed.), *Global Physical Climatology (Second Edition)* (pp. 49-94). Elsevier.  
<https://doi.org/10.1016/B978-0-12-328531-7.00003-7>